

# Recommended Childhood and Adolescent Immunization Schedule UNITED STATES • 2006

Vaccine ▼	Age ►	Birth	1 month	2 months	4 months	6 months	12 months	15 months	18 months	24 months	4–6 years	11–12 years	13–14 years	15 years	16–18 years
Hepatitis B <sup>1</sup>	HepB		HepB	HepB <sup>1</sup>		HepB						HepB Series			
Diphtheria, Tetanus, Pertussis <sup>2</sup>			DTaP	DTaP	DTaP			DTaP		DTaP	Tdap		Tdap		
<i>Haemophilus influenzae</i> type <sup>3</sup>			Hib	Hib	Hib <sup>2</sup>		Hib								
Inactivated Poliovirus			IPV	IPV		IPV				IPV					
Measles, Mumps, Rubella <sup>4</sup>							MMR			MMR			MMR		
Varicella <sup>5</sup>							Varicella						Varicella		
Meningococcal <sup>6</sup>												MCV4		MCV4	
Pneumococcal <sup>7</sup>			PCV	PCV	PCV		PCV			PCV		PPV			
Influenza <sup>8</sup>							Influenza (Yearly)					Influenza (Yearly)			
Hepatitis A <sup>9</sup>												HepA Series			

Vaccines within broken line are for selected populations

This schedule indicates the recommended ages for routine administration of currently licensed childhood vaccines, as of December 1, 2005, for children through age 18 years. Any dose not administered at the recommended age should be administered at any subsequent visit when indicated and feasible. ■ Indicates age groups that warrant special effort to administer those vaccines not previously administered. Additional vaccines may be licensed

any components of the combination are indicated and other components of the vaccine are not contraindicated and if approved by the Food and Drug Administration for that dose of the series. Providers should consult the respective ACIP statement for detailed recommendations. Clinically significant adverse events that follow immunization should be reported to the Vaccine Adverse Event Reporting System (VAERS). Guidance about how to obtain and