

For children approximately 2 months-8 months give Acetaminophen 160 mg/5 ml every 4 hours

FOR Fevers above 100.4(forehead) or 99.4(underarm/axillary)
For infants less than 2 months with a fever call doctor or go to ER

For children approximately 8 months and older *alternate*
Ibuprofen 100 mg/5 ml and Acetaminophen 160 mg/5 ml every 4 hours

**DO NOT GIVE ACETAMINOPHEN AND IBUPROFEN AT THE SAME TIME.
ALTERNATE MEANS TO GIVE THE CHILD ACETAMINOPHEN THEN 4
HOURS LATER GIVE THE CHILD IBUPROPHEN. THEN REPEAT THIS
PROCESS EVERY 4 HOURS.**

<u>Weight</u>	<u>ACETAMINOPHEN</u> TYLENOL, TRIAMINCIC FEVER REDUCER/Pediacare Infants/Childrens 160 mg/5 ml	<u>IBUPROFEN</u> ADVIL, MOTRIN Infants 50 mg/1.25 ml	<u>IBUPROFEN</u> ADVIL, MOTRIN Childrens 100 mg/5 ml
6-11 (0-3 mos)	1.25 ml every 4 hours	NONE	NONE
12-17 lbs (4-8 mos)	2.5 ml every 4 hours	NONE	NONE
18-23 lbs (9-23 mos)	3.75 ml every 4 hours	1.5 ml every 8 hours	3.75 ml every 8 hours
24-35 lbs (24-36mos)	5 ml every 4 hours	1.875 ml-2 ml every 8 hours	5 ml every 8 hours
36-45 lbs	7.5 ml every 4 hours		7.5 ml every 8 hours
46-59 lbs	8. ml every 4 hours		8. ml every 8 hours
60 lbs and over	10. ml every 4 hours		10. ml every 8 hours